

[Home](#) > [Navigating Cancer Care](#) > [How Cancer is Treated](#) > [Surgery](#) > [Understanding Colostomies](#)

Printed March 5, 2015 from <http://www.cancer.net/navigating-cancer-care/how-cancer-treated/surgery/understanding-colostomies>

Understanding Colostomies [1]

This section has been reviewed and approved by the [Cancer.Net Editorial Board](#) [2], 04/2014

Key Messages:

- A colostomy is a type of surgery that diverts waste from the large intestine to the outside of the body and can either be temporary or permanent.
- It may be needed when part of the large intestine is removed or damaged as a result of several medical conditions, including some types of cancer.
- After having a colostomy, your health care team will teach you to care for the opening made during the colostomy and empty the colostomy bag.
- After recovering from a colostomy, you should expect to continue to do most of the activities and wear the same type of clothing that you could before surgery because modern colostomy products are less noticeable than in the past.

A colostomy is a surgical opening called a stoma that connects the large intestine to the outside of your abdomen, providing a pathway for waste to exit the body. The waste is then collected in a pouch worn on the outside of your body. A colostomy is used when a part of the large intestine is surgically removed or when the flow of stool needs to be diverted from exiting via the rectum due to partial blockage or damage to the large bowel.

The large intestine, also called the large bowel, is made up of the colon and rectum and is connected to the small intestine. The small intestine, or the small bowel, is where nutrients—including proteins, fats, and carbohydrates—are digested and absorbed into blood vessels. Remaining food that cannot be digested moves from the small intestine to the large intestine, specifically to the colon, which absorbs water from the waste and stores the waste until the next bowel movement.

Several of the following medical conditions affect the colon so that part or all of it is unable to function as it should:

- An abdominal infection caused by a ruptured colon
- An externally caused injury to the colon or rectum
- [A blockage of the large intestine](#) [3]
- An abnormal connection between the anus and reproductive organs
- Certain types of cancer, such as [colorectal cancer](#) [4]

Most patients with cancer who need a colostomy only need it temporarily?usually for several months?to help the colon or the rectum heal after a portion has been removed. However, in some situations, a permanent colostomy may be needed.

Types of colostomy

There are several types of colostomies based on where in the four main sections of the colon it is needed.

Ascending colostomy. An ascending colostomy is located in the ascending colon, which extends from the beginning of the large intestine to the right side of the abdomen. In this procedure, only part of the colon remains functional. As a result, little water is absorbed from the waste, and the discharge is usually liquid. However, this type of procedure is rare because an ileostomy is more appropriate for the ascending colon. An ileostomy is a procedure in which the small intestine is diverted to the abdominal wall and waste is emptied into a pouch.

Transverse colostomy. A transverse colostomy is located in the transverse colon, which crosses the top of the abdomen. Transverse stool is usually soft or loose because only a small portion of the colon has absorbed water from the indigestible material. This is a common type of colostomy, and there are three versions of it. For patients with a loop colostomy, the colostomy stoma has one opening, and stool can exit through that opening. In this type, the colon is not disconnected, and patients will sometimes pass stool or gas through the rectum. For patients with a single-barrel colostomy, the colon?including the rectum and anal opening?is removed below the colostomy. This type of colostomy is permanent. For patients with a double-barrel colostomy, the colon is divided, and the two ends form two separate stomas. Stool exits from one of the stomas, while mucus made by the colon exits from the other. This type of colostomy is not as common as the other two types.

Descending colostomy. A descending colostomy is located in the descending colon, which takes waste down the left side of the abdomen. The stool there is usually firm because it has moved across functioning portions of the colon.

Sigmoid colostomy. A sigmoid colostomy?the most common type?is located in the sigmoid colon, the bottom portion of the large intestine, which moves waste to the rectum. Sigmoid colostomies produce stool that is more solid and regular than other colostomies.

The surgical procedure

The surgery to create a colostomy may be done through a large surgical incision or cut in the abdomen or through small incisions in the skin, called laparoscopic surgery [5]. Laparoscopic surgery is a minimally invasive type of surgery that minimizes pain and post-operative recovery time. With both types of surgery, the patient is given general anesthesia [6].

The surgeon stitches one end of the colon to a stoma. A special bag is then attached to the stoma, allowing solid waste and gas from the colon to empty into it. Physicians usually place stomas on a smooth skin surface on the stomach in an area where the patient can easily see it and take care of it.

Risks associated with the surgery include bleeding inside the colon, damage to nearby organs, infection, and bleeding from the stoma.

Recovery from surgery

A complete recovery after receiving a colostomy may take up to two months, with most patients needing to stay in the hospital for up to one week after the procedure. Eating is restricted during this time to help the colon heal. If the colostomy is temporary, a reversal surgery may be performed approximately 12 weeks later, after the colon has healed.

Colostomy care

Once you have recovered from surgery, you will need to empty the colostomy bag several times per day, preferably when the bag is less than one-half full. There are several types of colostomy pouches available, including open-end or drainable and closed-end or disposable pouches. Some pouches have a filter that deodorizes and vents gas so that the bag does not become too stretched and come off of the abdomen or burst.

It is important to keep the skin surrounding the stoma, called peristomal skin, clean to avoid infection. The stoma will always appear red and may bleed occasionally, which is normal. However, bleeding should not continue for long periods.

Amounts of stool and gas deposited into the pouch will vary based on the type of colostomy you received and your diet. Your doctor or nurse can provide information on the types of foods that you should avoid to reduce gas.

You will not be able to control when stool and gas move into the pouch because the colon cannot voluntarily block and hold waste like the rectum does.

Living with a colostomy

Although a colostomy will alter your life, modern colostomy products are designed to be less inconvenient and allow you to do most of the activities that you could before surgery. You should be able to wear the same type of clothes you wore before the colostomy because pouches are designed to blend in, fitting close to the body. Products also have odor-barrier film that traps bad odor, which only releases when the colostomy pouch is being emptied.

A colostomy should not restrict you from the activities of your daily life. Patients with colostomies go to work, play most sports, and engage in sexual activity [7]. Reasonable activity will not hurt you or your stoma.

Despite efforts to adjust, some people may feel depressed or embarrassed by their colostomy. Reaching out to a support group [8] of patients with colostomies, either in person or online, may

help answer your questions and provide emotional support. Learn more about [body image and cancer](#) [9].

More Information

[Understanding Cancer Surgery](#) [5]

Additional Resource

[MedlinePlus: Colostomy](#) [10]

Links:

[1] <http://www.cancer.net/navigating-cancer-care/how-cancer-treated/surgery/understanding-colostomies>

[2] <http://www.cancer.net/about-us>

[3] <http://www.cancer.net/node/25244>

[4] <http://www.cancer.net/node/18701>

[5] <http://www.cancer.net/node/24720>

[6] <http://www.cancer.net/node/24375>

[7] <http://www.cancer.net/node/25240>

[8] <http://www.cancer.net/node/25383>

[9] <http://www.cancer.net/node/25194>

[10] <http://www.nlm.nih.gov/medlineplus/ency/article/002942.htm>